

Platynotini (Coleoptera, Tenebrionidae, Opatrinae) of the Natural History Museum in Geneva

by

Dariusz IWAN *

With 7 figures

ABSTRACT

Platynotini (Coleoptera, Tenebrionidae, Opatrinae) of the Natural History Museum in Geneva. — 29 species of Platynotini are recorded from the Geneva collection, and new faunistic records are provided. One new species — *Platydendarus marzenae* is described and figured.

Through the kindness of Dr. Ivan Löbl of the Natural History Museum in Geneva I was able to examine the interesting collection of the Platynotini. The group is poorly known and only few faunistic and taxonomic papers were published (MULSANT and REY, 1853; GRIDELLI, 1947; KOCH, 1956; KASZAB, 1975).

The mentioned material is deposited in the Natural History Museum in Geneva.

I thank Dr. Ivan Löbl and Dr. Ottó Merkl (Budapest) for the loan of material and Dr. Stanisław A. Słipiński for his help and advice.

Opatriinus aciculatus Le Conte

Opatriinus aciculatus Le Conte, 1858, Proc. Ac. Phil. 1858: 75; Horn 1870, Trans. Am. phil. Soc. 14: 349.

MATERIAL EXAMINED: "Texas", 1♀.

Distribution: USA: Texas.

* Institute of Zoology, Wilcza 64, 00-679 Warszawa, Poland.

Opatriinus gemellatus (Olivier)

Blaps gemellata Olivier, 1795, Ent. 3, 60: 9. — In *Opatriinus*: Mulsant and Rey 1853: 299; Marcuzzi 1977, *Studies Curaçao* 12, 170: 22.

MATERIAL EXAMINED: “Guadeloupe s. str., Deshaies Gr. Anse, 26.VII.1974, rotten leaves”, (Vit), 9♀ 9♂ 8♂ ♂. — “Guadeloupe Grd. T., St. Louis, 21.XI.1983”, (Vit), 2♀ 9♂ 1♂. — “Venezuela, coll. Gory”, 2♀ 9♂ 2♂ ♂. — “Colombia”, 2♂ ♂ 1♀. — “Porto Cabello”, 1♀.

Distribution: Guiana, Venezuela, Colombia, Margarita, Trinidad, Tobago, Grenada, Bequia, Moustique, Union, Barbados, St. Vincent, St. Lucia, Martinique, Dominica, Les Saintes, Marie-Galante, Guadeloupe, St. Kitts, St. Croix.

Opatriinus laticollis Latreille

Opatriinus laticollis Latreille, 1833, *Voy. Humb. Bonpl.* 34: 66; Mulsant and Rey 1853: 301.

MATERIAL EXAMINED: “Colombia”, 1♀ 1♂.

Distribution: New Grenada, Colombia.

Opatriinus minimus (Beauvois)

Tenebrio minimus Beauvois, 1805, *Ins. Afr. Amer.* 31: 164. — In *Opatriinus*: Chevrolat 1852, *Ann. Fr.* 2, 10: 636;

Opatrium notum Say, 1827, *Journ. Ac. Phil.* 5: 237; Mulsant and Rey, 1853: 309.

MATERIAL EXAMINED: “Florida, Wakolla Springs bei Tallahassee”, (Smetana), 1♀. — “Florida, Lake Istokpoga near Lorida, 12.III.1968”, (Smetana), 2♂ ♂. — “Louisiana, St. John The Baptit Co., Manchac, 20.III.1968”, (Smetana), 1♀. — “North Amer., coll. Gory”, 1♀ 3♂ ♂. — “Louisiana”, 1♀.

Distribution: Atlantic coast of the USA from Louisiana to Massachusetts.

Opatriinus moestus Mulsant and Rey

Opatriinus moestus Mulsant and Rey, 1853: 306.

MATERIAL EXAMINED: “Argentina, Buenos Aires, Stevenia, XI.1950”, (Mussard), 1♀.

Distribution: Argentina, Brazil.

Opatriinus pullus (Sahlberg)

Opatrium pullus Sahlberg, 1823, *Thon's Arch.* II: 16. — In *Opatriinus*: Gebien 1910, *Coleopterorum Catalogus* 15: 277; Marcuzzi 1977, *Studies Curaçao* 12, 170: 32; *Opatriinus anthracinus* Mulsant and Rey, 1853: 304.

MATERIAL EXAMINED: “Cuba”, 2♀ 9♂ 1♂. — “Mexico, Merida, Yucatan, 13-15.XII.1975”, (de Chambrier), 3♀ 9♂ 1♂.

Distribution: Puerto Rico, Jamaica, Cuba, Mexico.

Opatriinus corvinus Mulsant and Rey

Opatriinus corvinus Mulsant and Rey, 1853: 296; Gridelli 1947: 40; Koch 1956: 94.

MATERIAL EXAMINED: “Egypt”, (Melly), 2♀ 9♂ 2♂ ♂.

Distribution: Mali, Niger, Chad, Sudan, Ethiopia, Egypt.

Opatriinus costulatus (Guérin)

Opatrium costulatum Guérin, 1849, *Voy. Lefebure* 5: 321. — In *Opatriinus*: Gemminger and Harold 1870, *Catalogus Coleopterorum* VII: 1915; Gridelli 1947: 48; Koch 1956: 110.

MATERIAL EXAMINED: "Abyssinia", 2♂♂ 4♀♀.

Distribution: Nigeria, Chad, Sudan, Ethiopia, Somaliland.

Opatriinus latipes (Sahlberg)

Opatrium latipes Sahlberg, 1823, *Thon's Arch.* II, 1: 13. — In *Opatriinus*: Gemminger and Harold 1870, *Catalogus Coleopterorum* VII: 1915; Gridelli 1947: 46; Koch 1956: 105;

Opatriinus ovalis Mulsant and Rey, 1853: 315;

Opatriinus atratus Quedenfeldt, 1885, *Berl. Ent. Zeitchr.* XXIX: 8;

Opatriinus opacus Gebien, 1904, *Ark. Zool.* 2, 5: 4.

MATERIAL EXAMINED: "Ivory Coast, Adzopé", (Mussard), 1♂. — "Ivory Coast, Sassandra, August 1948", (Mussard), 2♀♀. — "Cameroon, at light", (Notari), 2♂♂ 1♀. — "Upper Volta, Ouagadougou, V-VII.1974", (Roler), 1♂. — "Senegal, coll. Gory", 1♀. — "Congo, Luluabourg", 2♀♀.

Distribution: tropical Africa.

Opatriinus servus Mulsant and Rey

Opatriinus servus Mulsant and Rey, 1853: 317; Gridelli 1947: 92; Koch 1956: 112.

MATERIAL EXAMINED: "Cameroon, VII.1983, Etoubi Assok, at light", (Notari), 1♂.

Distribution: tropical Africa.

Anchophthalmus algoensis Péringuay

Anchophthalmus algoensis Péringuay, 1904, *Ann. S. Afr.* 3: 237; Koch 1956: 128.

MATERIAL EXAMINED: "South Africa: Transvaal", (Audéoud), 1♂. — "Mozambique, Lourenço Marques", (Audéoud), 1♂.

Distribution: southern Mozambique, northern Transvaal.

Anchophthalmus plicipennis Péringuay

Anchophthalmus plicipennis Péringuay, 1904, *Ann. S. Afr.* 3: 238; Koch 1956: 148.

MATERIAL EXAMINED: "Tanganyika, Shinyanga", 1♂ 3♀♀.

Distribution: south-eastern part of tropical Africa.

Glyptopteryx quadricollis (Fairmaire)

Selinus quadricollis Fairmaire, 1887, *Ann. Soc. Ent. France* 6: 284. — In *Glyptopteryx*: Koch 1956: 217;

Glyptopteryx forticostis Gebien, 1910, *Erg. Exp. Kilim.* 7: 376.

MATERIAL EXAMINED: "Kenya, lac Naivasha, 10 km Fisherman's Camp, 5.XI.1974", (Mahnert, Perret), 1♀.

Distribution: eastern part of tropical Africa.

Melanopterus porcatus Mulsant and Rey

Melanopterus porcatus Mulsant and Rey, 1854, *Mem. Ac. Lyon* 4: 159.

MATERIAL EXAMINED: "Cape Province", 1♀.

Distribution: South Africa: Cape Province.

Monodius convexipennis (Gebien)

Selinus convexipennis Gebien, 1904, *Ark. Zool.* 5: 2. — In *Monodius*: Koch 1956: 181.

MATERIAL EXAMINED: "Cameroon, Etoubi Assok, at light, VII.1983", (Notari), 3♀♀ 1♂.

Distribution: Nigeria, Cameroon.

Quadrideres femineus (Lesne)

Selinus femineus Lesne, 1922, *Voy. Roths. 1: 701.* — In *Quadrideres*: Koch 1956: 192.

MATERIAL EXAMINED: "Kenya, lac Nivasha, 10 km Fisherman's Camp, 5.XI.1974", (Mahnert, Perret), 7♀♀ 3♂♂.

Distribution: eastern part of tropical Africa.

Quadrideres simplicipes (Gebien)

Selinus simplicipes Gebien, 1910, *Erg. Exp. Kilim.* 7: 374. — In *Quadrideres*: Koch 1956: 205.

MATERIAL EXAMINED: "Kenya, Taita Hills, Mwarungu, 1700 m, 3.II.1974", (Mahnert, Perret), 3♀♀.

Distribution: Kenya.

Selinus planus (Fabricius)

Opatrium planum Fabricius, 1792, *Ent. Syst. I:* 118. — In *Selinus*: Mulsant and Rey 1853: 324; Koch 1956: 251.

MATERIAL EXAMINED: "West Africa", 1♀.

Distribution: north-western part of tropical Africa.

Eucolus polinieri Mulsant and Rey

Eucolus Polinieri Mulsant and Rey, 1853: 292; Kaszab 1975: 285.

MATERIAL EXAMINED: "Ceylon, Kandy", 1♀ 1♂.

Distribution: southern India, Sri Lanka.

Menearchus dispar (Herbst)

Tenebrio dispar Herbst, 1797, *NaturSystem*, 7: 248. — In *Menearchus*: Kaszab 1975: 336.

MATERIAL EXAMINED: "India or.", 2♂♂. — "India mér. 24.V.". (Carl et Escher), 1♀. — "Ceylon, Tricomalie" (Humbert), 3♀♀ 2♂♂.

Distribution: southern India, Sri Lanka.

Menearchus hirtipes Kaszab

Menearchus hirtipes Kaszab, 1975: 335.

MATERIAL EXAMINED: "India, Madras, Coimbatore, 1400 ft", (Nathan), 1♂ 3♀ ♀.

Distribution: southern India.

Menearchus laevipennis Kaszab

Menearchus laevipennis Kaszab, 1975: 347.

MATERIAL EXAMINED: "Ceylon, Trincomalie" (Humbert), 4♂ ♂.

Distribution: Sri Lanka.

Menearchus tenuestriatus (Fairmaire)

Pseudoblaps tenuestriata Fairmaire, 1898, *Annls Soc. ent. Fr.* 67: 391. — In *Menearchus*: Kaszab 1975: 342.

MATERIAL EXAMINED: "Ceylon, Trincomalie" (Humbert), 2♂ ♂ 3♀ ♀.

Distribution: Sri Lanka.

Platydendarus javanus (Wiedemann)

Opatrium javanum Wiedemann, 1819, *Zool. Mag.* 1, 3: 163. — In *Platydendarus*: Kaszab 1975: 318.

MATERIAL EXAMINED: "Thailand, Huà Dông, 20 km N Nakhon Sawan, 1981", (Schwendinger), 1♀ 1♂. — "Thailand, Surin, 27.XII.1985", (Schwendinger), 1♀. — "Thailand, Chiang Mai", (Schwendinger), 3♀ ♀ 3♂ ♂.

Distribution: southern China, India, Sunda Islands, Moluccas.

Platydendarus marzenae n. sp. (figs 1-7)

Body black, shiny. Head densely and coarsely punctured; frons strongly depressed medially and near fronto-clypeal suture (fig. 2); eyes convex. Pronotum (fig. 3) $0,71 \times$ as long as wide; disc densely and coarsely punctured, wrinkled towards lateral margins; sides of pronotum distinctly rounded, with shallow sinuosity in front of posterior angles; pronotal edges crenulate; anterior angles acute, slightly prominent; pronotal base sinuate laterally, marginal carina obsolescent and broadly interrupted medially. Elytra (fig. 4) $1,4 \times$ as long as wide and $2,4 \times$ as long as pronotum, broadly oval, convex laterally, with rounded sides; elytral punctures strong, round, situated in distinct grooves; fourth row consist of 36 punctures; intervals slightly convex, with sparser, much finer punctuation. External female genitalia as on figure 7; total length 1,92 mm, paraprocts 1,06 mm, coxites 1,12 mm, ratio p/c 0,96; paraprocts trapezoidal, with longitudinal baculi; coxites subdivided into 4 lobes; gonostyli with a number of hairs at the top, one of them longer than the other. Bursa copulatrix (figs 5, 6) with three well-sclerotised membranes.

Length 12,5-13,1 mm.

HOLOTYPE ♀ — [Burma, Taninthari], "India or., Tenasserim, Meešan Fea. Apr. 1887, It. L. Fea Dupl. Ach. 7.92", (Mus. Geneva).

PARATYPE ♀ — [Burma, Taninthari], "Tenasserim, J. W. M.", (Hungarian Natural History Museum).

Dedicated to Miss Marzena Kowalow.


This species is similar to *P. rondoni* Kaszab, 1975. It differs from it in having the lateral edges of pronotum crenulate (smooth in *rondoni*) and elytral punctures in distinct grooves (without grooves in *rondoni*).



1

FIG. 1.

Platydendarus marzenae sp. n., dorsal view.



FIGS 2-7.

Platydendarus marzenae sp. n. — 2: head, dorsal; 3: pronotum; 4: elytra; 5: bursa copulatrix, dorsal; 6: bursa copulatrix, lateral; 7: ovipositor, dorsal.

Platynotooides belli (Fairmaire)

Platynotus Belli Fairmaire, 1896, *Annls. Soc. ent. Belg.* 40: 13. — In *Platynotooides*: Kaszab 1975: 297.

MATERIAL EXAMINED: "India", 2♀♀.

Distribution: southern India, Burma.

Rugoplatynotus andrewesi (Fairmaire)

Pseudoblaps Andrewesii Fairmaire, 1896, *Annls. Soc. ent. Belg.* 40: 14. — In *Rugoplatynotus*: Kaszab 1975: 289.

MATERIAL EXAMINED: "Burma, Thigyara, 11.VI.1885", (Fea), 1♀.

Distribution: Burma.

Notocorax mandli Kaszab

Notocorax mandli Kaszab, 1975: 302.

MATERIAL EXAMINED: "India, Madras, Coimbatore, 1400 Ft, VIII.1971", (Nathan), 1♂. — "India, Madras, Coimbatore, sub lapido, oct. 1976", 1♀.

Distribution: southern India.

Notocorax nervosus Mulsant and Rey

Notocorax nervosus Mulsant and Rey, 1853: 50.

MATERIAL EXAMINED: "India, Madras, Coimbatore, 1400 Ft, IX.1972", (Nathan), 2♂♂. — "India, Madras, Coimbatore, 1400 Ft, XI.1975", (Nathan), 1♀.

Distribution: southern India.

BIBLIOGRAPHY

- GRIDELLI, E. 1947. Specie africane del genere *Opatrinus* Muls. Rey (Coleopt. Tenebr.). *Atti Mus. civ. Stor. nat. Trieste* 16, 3: 37-52.
- KASZAB, Z. 1975. Revision der Asiatischen Platynotinen (Coleoptera: Tenebrionidae). *Acta zool. hung.* 21, 3-4: 277-367.
- KOCH, C. 1956. Il Tenebrionidae (Coleoptera Polyphaga) Opatrinae, First part: Platynotini, Litoborini and Loensini. Exploration du Parc National de l'Upemba, *Inst. Parcs Nat. Congo Belge, Bruxelles*, 1-472 pp., 8 maps, 282 figs, 35 pl.
- MULSANT, E. and Cl. REY. 1853. Essai d'une division des derniers Mélasomes. *Opusc. ent.* 4: 1-242.